

A NOTE ON THE WEAK LEFSCHETZ PROPERTY OF MONOMIAL COMPLETE INTERSECTIONS IN POSITIVE CHARACTERISTIC

HOLGER BRENNER AND ALMAR KAID

ABSTRACT. Let K be an algebraically closed field of characteristic $p > 0$. We apply a theorem of C. Han to give an explicit description for the weak Lefschetz property of the monomial Artinian complete intersection $A = K[X, Y, Z]/(X^d, Y^d, Z^d)$ in terms of d and p . This answers a question of J. Migliore, R. M. Miró-Roig and U. Nagel and, equivalently, characterizes for which characteristics the rank-2 syzygy bundle $\text{Syz}(X^d, Y^d, Z^d)$ on $\mathbb{P}^2$ satisfies the Grauert-Mülich theorem. As a corollary we obtain that for $p = 2$ the algebra A has the weak Lefschetz property if and only if $d = \lfloor \frac{2^t+1}{3} \rfloor$ for some positive integer t . This was recently conjectured by J. Li and F. Zanello.

Mathematical Subject Classification (2010): primary: 13D02, 13E10, 14J60, secondary: 13C13, 13C40, 14F05.

Keywords: syzygy, stable bundle, Grauert-Mülich Theorem, weak Lefschetz property, Artinian algebra, monomial complete intersection.

1. INTRODUCTION

Let $R = K[X_0, \dots, X_N]$ be the polynomial ring in $N + 1$ variables over an algebraically closed field K and let $f_1, \dots, f_n$ denote R_+ -primary homogeneous polynomials in R (i.e., $\sqrt{(f_1, \dots, f_n)} = R_+$). Then the quotient $A := R/(f_1, \dots, f_n)$ is an Artinian graded K -algebra, i.e., A is of the form

$$A = K \oplus A_1 \oplus \dots \oplus A_s$$

for some integer $s \geq 0$. The algebra A has the *weak Lefschetz property* (abbreviated by WLP) if for every general linear form $\ell \in R_1$ the multiplication maps

$$A_m \xrightarrow{\cdot \ell} A_{m+1}$$

have maximal rank for $m = 0, \dots, s - 1$.

We also associate to the polynomials $f_1, \dots, f_n$ the *syzygy bundle* on $\mathbb{P}^N = \text{Proj } R$. This vector bundle is given by the short exact sequence

$$0 \longrightarrow \text{Syz}(f_1, \dots, f_n) \longrightarrow \bigoplus_{i=1}^n \mathcal{O}_{\mathbb{P}^N}(-d_i) \longrightarrow \mathcal{O}_{\mathbb{P}^N} \longrightarrow 0,$$

where $d_i := \deg(f_i)$. If $N = 2$ and $\text{char}(K) = 0$, we gave in our article [2] a characterization for the weak Lefschetz property of the Artinian algebra A in terms of the generic splitting type of the syzygy bundle $\text{Syz}(f_1, \dots, f_n)$ (see [2, Theorem 2.2]). As a consequence we obtained, using the theorem of Grauert-Mülich (see [6, Theorem 3.0.1]), the result of Harima-Migliore-Nagel-Watanabe saying that *every* Artinian complete intersection in $K[X, Y, Z]$ has the weak Lefschetz property (see [5, Theorem 2.4] and [2, Corollary 2.4]). The easy examples of the stable syzygy bundles $\mathcal{S} = \text{Syz}(X^p, Y^p, Z^p)$ over a field of characteristic p show that neither Grauert-Mülich ($\mathcal{S}$ splits on every line $L \subset \mathbb{P}^2$ as $\mathcal{S}|_L \cong \mathcal{O}_L(-p) \oplus \mathcal{O}_L(-2p)$; cf. also the example of L. Ein in [3, Section 4]) nor the theorem of Harima et al. holds in positive characteristic (cf. [8, Example 7.10]).

The aim of this paper is to give a numerical characterization of the WLP for monomial Artinian complete intersections $K[X, Y, Z]/(X^d, Y^d, Z^d)$ in positive characteristic. This answers [8, Question 7.12] of Migliore-Miró-Roig-Nagel and, equivalently, characterizes for which characteristics the rank-2 syzygy bundle $\text{Syz}(X^d, Y^d, Z^d)$ on $\mathbb{P}^2$ satisfies the Grauert-Mülich theorem. As a consequence we obtain a proof for the recent conjecture [7, Conjecture 3.9] of J. Li and F. Zanello.

Besides our geometric approach, the key ingredient for our investigation is a theorem of C. Han which computes the *syzygy gap* for an ideal of the form $(X^d, Y^d, (X + Y)^d)$ in $K[X, Y]$.

2. A CHARACTERIZATION OF THE WLP FOR MONOMIAL ARTINIAN COMPLETE INTERSECTIONS IN POSITIVE CHARACTERISTIC

The following lemma relates the question of whether the Artinian algebra $A := K[X, Y, Z]/(X^d, Y^d, Z^d)$ has the weak Lefschetz property to the generic splitting type of the corresponding syzygy bundle $\text{Syz}(X^d, Y^d, Z^d)$ on the projective plane à la Grauert-Mülich.

Lemma 2.1. *Let K be an algebraically closed field. Then the Artinian complete intersection $K[X, Y, Z]/(X^d, Y^d, Z^d)$ has the weak Lefschetz property if and only if the syzygy bundle $\mathcal{S} := \text{Syz}(X^d, Y^d, Z^d)$ on $\mathbb{P}^2$ splits on a generic line L as $\mathcal{S}|_L \cong \mathcal{O}_L(a) \oplus \mathcal{O}_L(b)$ with $a \geq b$ and $0 \leq a - b \leq 1$.*

Proof. Since $\text{Syz}(X^d, Y^d, Z^d)$ is a stable vector bundle (see for instance [1, Corollary 3.2]), the assertion of the lemma follows from [2, Theorem 2.2] (note that the proof of this result holds in any characteristic). $\square$

Next, we explain the notion of *syzygy gap* introduced in [9] by P. Monsky. Let K be an algebraically closed field and consider the ideal $I := (X^{d_1}, Y^{d_2}, (X + Y)^{d_3})$ in $S := K[X, Y]$. The minimal graded free resolution of the quotient S/I is given by

$$0 \longrightarrow S(a) \oplus S(b) \longrightarrow S(-d_1) \oplus S(-d_2) \oplus S(-d_3) \longrightarrow S \longrightarrow S/I \longrightarrow 0,$$

with integers a, b , $a \geq b$. The difference $\delta(d_1, d_2, d_3) := a - b$ is called the *syzygy gap* and constitutes a function $\delta : \mathbb{N}^3 \rightarrow \mathbb{N}$. It is easy to see that $a + b = -(d_1 + d_2 + d_3)$ and hence $\delta(d_1, d_2, d_3) \equiv d_1 + d_2 + d_3 \pmod{2}$.

Corollary 2.2. *Let K be an algebraically closed field (of any characteristic), $A = K[X, Y, Z]/(X^d, Y^d, Z^d)$ and denote by $\mathcal{S} = \text{Syz}(X^d, Y^d, Z^d)$ the corresponding syzygy bundle. Then the following conditions are equivalent.*

- (1) *The algebra A has the weak Lefschetz property.*
- (2) *The bundle $\mathcal{S}$ splits on a generic line L as $\mathcal{S}|_L \cong \mathcal{O}_L(a) \oplus \mathcal{O}_L(b)$ with $a \geq b$ and $0 \leq a - b \leq 1$ (i.e., the theorem of Grauert-Mülich holds).*
- (3) *We have $\delta(d, d, d) \leq 1$.*

Proof. The equivalence (1) $\Leftrightarrow$ (2) is proved in Lemma 2.1.

(2) $\Leftrightarrow$ (3). If we want to compute the splitting type of $\mathcal{S}$ on a line L given by the equation $Z = uX + vY$ with coefficients $u, v \in K$, $u, v \neq 0$ (in particular this holds for a generic line), we can assume without loss of generality that $u = v = 1$. Hence computing the generic splitting type of $\mathcal{S}$ is the same as computing the syzygy gap $\delta(d, d, d) = a - b$. $\square$

We denote by $\delta^* : [0, \infty)^3 \rightarrow [0, \infty)$ the continuous continuation of δ ; see [9, Definition 19 and the following] for this function and some of its properties. We set

$$L_{\text{odd}} := \{(u_1, u_2, u_3) \in \mathbb{Z}^3 : \sum_{i=1}^3 u_i \text{ odd}\} \subset \mathbb{Z}^3.$$

An element $u = (u_1, u_2, u_3) \in \mathbb{Z}^3$ belongs to L_{odd} if and only if all entries of u are odd or if there is only one odd entry u_i , $i \in \{1, 2, 3\}$. Further, we denote by td the *taxi-cab distance* in $\mathbb{R}^3$ defined as $\text{td}(v, w) := \sum_{i=1}^3 |v_i - w_i|$ for triples $v = (v_1, v_2, v_3), w = (w_1, w_2, w_3) \in \mathbb{R}^3$.

The following theorem due to C. Han yields an effective way to compute δ^* for a given triple $v = (v_1, v_2, v_3) \in [0, \infty)^3$.

Theorem 2.3 (Han). *Let K be an algebraically closed field of characteristic $p > 0$ and assume the entries of $v = (v_1, v_2, v_3) \in [0, \infty)^3$ satisfy $v_1 \leq v_2 \leq v_3$ and $v_3 < v_1 + v_2$. If there exists $s \in \mathbb{Z}$ and a triple $u = (u_1, u_2, u_3) \in L_{\text{odd}}$ such that $m := \text{td}(p^s v, u) < 1$, then there exists such a pair s, u with minimal s . With these data s, u and m we have*

$$\delta^*(v) = p^{-s}(1 - m).$$

If no such pair exists, then $\delta^(v) = 0$.*

Proof. See [4, Theorems 2.25 and 2.29] or [9, Corollary 23] for an easier proof. $\square$

Lemma 2.4. *Let $d \in \mathbb{N}_+$ and p be a prime number. Then the following conditions are equivalent.*

(1) *There exists $k \in \mathbb{N}$ and $n \in \mathbb{N}$ such that*

$$\frac{3d}{6k+2} > p^n > \frac{3d}{6k+4}.$$

(2) *There exists an odd number $u \in \mathbb{N}$ and $s \in \mathbb{Z}$, $s \leq 0$, such that*

$$u - \frac{1}{3} < dp^s < u + \frac{1}{3}.$$

(3) *There exists an integer s , $s \leq 0$, such that the taxi-cab distance of (dp^s, dp^s, dp^s) to some point in L_{odd} is < 1 .*

Proof. To proof the equivalence between (1) and (2) we set $s = -n$ and $u = 2k + 1$. The condition in (1) is equivalent with

$$\frac{3}{3u-1} > \frac{p^n}{d} > \frac{3}{3u+1}$$

and by inverting it is equivalent with

$$u - \frac{1}{3} < dp^s < u + \frac{1}{3}.$$

If (2) is true, then we have $(u, u, u) \in L_{\text{odd}}$ and the taxi-cab distance between (dp^s, dp^s, dp^s) and (u, u, u) is < 1 . On the other hand, the distance of a point on the diagonal to any point in L_{odd} outside the diagonal is at least 1, so we only have to consider points on the diagonal. $\square$

Lemma 2.5. *Let $d \in \mathbb{N}_+$ and p be a prime number. Suppose that there exists $0 \leq n' < n$ and $k', k \in \mathbb{N}$ such that*

$$\frac{3d-1}{6k'+2} > p^{n'} > \frac{3d+1}{6k'+4}$$

and

$$\frac{3d}{6k+2} > p^n > \frac{3d}{6k+4}.$$

Then

$$\frac{3d-1}{6k+2} > p^n > \frac{3d+1}{6k+4}.$$

Proof. Otherwise we would have either

$$\frac{3d+1}{6k+4} \geq p^n > \frac{3d}{6k+4}$$

or

$$\frac{3d}{6k+2} > p^n \geq \frac{3d-1}{6k+2}.$$

This gives either

$$p^n(6k+4) = 3d+1$$

or

$$p^n(6k+2) = 3d-1.$$

We plug this in the first inequality and get in the first case

$$\frac{p^n(6k+4)-2}{6k'+2} > p^{n'} > \frac{p^n(6k+4)}{6k'+4}$$

and by dividing through p^n we get

$$\frac{3k+2-\frac{1}{p^n}}{3k'+1} > p^{n'-n} > \frac{3k+2}{3k'+2}.$$

By inverting we obtain

$$\frac{3k'+1}{3k+2-\frac{1}{p^n}} < p^{n-n'} < \frac{3k'+2}{3k+2}.$$

From the right hand side we get $p^{n-n'} \leq \frac{3k'+1}{3k+2}$ which yields the contradiction

$$\frac{3k'+1}{3k+2-\frac{1}{p^n}} < \frac{3k'+1}{3k+2}.$$

In the second case we obtain

$$\frac{p^n(6k+2)}{6k'+2} > p^{n'} > \frac{p^n(6k+2)+2}{6k'+4}$$

and similar manipulations yield a contradiction. $\square$

The following theorem gives an explicit answer to [8, Question 7.12]. This question was also answered in [7, Corollary 3.6] but in a less explicit way.

Theorem 2.6. *Let K be a field of characteristic $p > 0$ and consider the monomial Artinian complete intersection $A := K[X, Y, Z]/(X^d, Y^d, Z^d)$. Then the following holds:*

- (1) *If d is even, then A does not have the weak Lefschetz property if and only if there exists a $k \in \mathbb{N}$ and an $n \in \mathbb{N}_+$ such that*

$$\frac{3d}{6k+2} > p^n > \frac{3d}{6k+4}.$$

- (2) *If d is odd, then A does not have the weak Lefschetz property if and only if there exists a $k \in \mathbb{N}$ and an $n \in \mathbb{N}_+$ such that*

$$\frac{3d-1}{6k+2} > p^n > \frac{3d+1}{6k+4}.$$

Proof. We prove (1). Assume that we have

$$\frac{3d}{6k+2} > p^n > \frac{3d}{6k+4}$$

for some $k \in \mathbb{N}$ and $n \in \mathbb{N}_+$. We set $s := -n$, $u := 2k+1$. Then we have $m := \text{td}(p^s(d, d, d), (u, u, u)) < 1$ by Lemma 2.4 and hence $\delta^*(d, d, d) = p^{-s}(1-m) > 0$. Since $\delta^*(d, d, d) = a - b$ and $a + b = -3d$ we must have $\delta^*(d, d, d) \geq 2$. We apply Corollary 2.2 and see that A does not have the WLP.

Now we assume that the numerical condition does not hold. Then by Lemma 2.4 there is no $s \leq 0$ such that the taxi-cab distance from $p^s(d, d, d)$ to an element $(u, u, u) \in L_{\text{odd}}$ is < 1 . This is also true for $s > 0$ since d is even. Hence it follows from Han's Theorem 2.3 that $\delta^*(d, d, d) = 0$ which implies by Corollary 2.2 the WLP for the algebra A .

Next we prove (2). First we remark that, since d is odd, the condition

$$\frac{3d}{6k+2} > p^n > \frac{3d}{6k+4}$$

is always fulfilled for $n = 0$ and k such that $d = 2k + 1$. We choose $n > 0$ maximal such that

$$\frac{3d}{6k+2} > p^n > \frac{3d}{6k+4}$$

holds for some k . Hence we can apply Han's Theorem 2.3 with $s := -n$ (minimal) and $u := 2k + 1$ to compute the syzygy gap.

Suppose that the numerical condition of part (2) is fulfilled for some $k' \in \mathbb{N}$ and $n' \in \mathbb{N}_+$. According to Lemma 2.5 we may assume that this condition also holds for the chosen (maximal) n , hence

$$\frac{3d-1}{6k+2} > p^n > \frac{3d+1}{6k+4}.$$

Then we have in particular

$$u - \frac{1}{3} = \frac{6k+2}{3} < dp^s < \frac{6k+4}{3} = u + \frac{1}{3}$$

by Lemma 2.4. Now we distinguish two cases.

Case 1: Let $u > dp^s$. Then the taxi-cab distance from $p^s(d, d, d)$ to the element $(u, u, u) \in L_{\text{odd}}$ equals

$$m := \text{td}(p^s(d, d, d), (u, u, u)) = 3(u - dp^s)$$

and we have $m < 1$ (by Lemma 2.4). So we obtain for the syzygy gap:

$$\begin{aligned} \delta^*(d, d, d) &= p^{-s}(1 - m) \\ &= p^{-s}(1 - 3u + 3dp^s) \\ &= p^{-s}(1 - 3u) + 3d \\ &= -p^n(6k+2) + 3d \\ &> -(3d-1) + 3d \\ &= 1. \end{aligned}$$

Therefore the syzygy gap is indeed ≥ 3 . Hence it follows from Corollary 2.2 that A does not have the WLP.

Case 2: Let $u \leq dp^s$. Then we obtain

$$m := \text{td}(p^s(d, d, d), (u, u, u)) = 3(dp^s - u)$$

which is again < 1 . So we can estimate the syzygy gap as follows:

$$\begin{aligned}
\delta^*(d, d, d) &= p^{-s}(1 - m) \\
&= p^{-s}(1 + 3u - 3dp^s) \\
&= (1 + 3u)p^{-s} - 3d \\
&= (6k + 4)p^n - 3d \\
&> 3d + 1 - 3d \\
&= 1.
\end{aligned}$$

Again we conclude that A does not have the WLP.

Next suppose that the numerical condition of part (2) does not hold. Then we have either

$$\frac{3d+1}{6k+4} \geq p^n > \frac{3d}{6k+4} \text{ or } \frac{3d}{6k+2} > p^n \geq \frac{3d-1}{6k+2},$$

where n and k are chosen as in the beginning of the proof of part (2).

Case 1: Let $\frac{3d+1}{6k+4} \geq p^n > \frac{3d}{6k+4}$. Then we even have

$$p^n(6k+4) = p^n(3u+1) = 3d+1.$$

Since

$$\frac{d}{u} = \frac{3d}{3u} > \frac{3d+1}{3u+1} = \frac{3d+1}{6k+4} = p^n,$$

we have $dp^s > u$. So we obtain

$$m := \text{td}(p^s(d, d, d), (u, u, u)) = 3(dp^s - u)$$

which is < 1 . This gives:

$$\begin{aligned}
\delta^*(d, d, d) &= p^{-s}(1 - m) \\
&= p^{-s}(1 + 3u - 3dp^s) \\
&= p^{-s}(1 + 3u) - 3d \\
&= p^n(3u+1) - 3d \\
&= 1.
\end{aligned}$$

Hence A has the WLP by Corollary 2.2.

Case 2: Let $\frac{3d}{6k+2} > p^n \geq \frac{3d-1}{6k+2}$. This implies

$$p^n(6k+2) = -p^n(1-3u) = 3d-1.$$

Since

$$p^n = \frac{3d-1}{6k+2} = \frac{3d-1}{3u-1} < \frac{d}{u},$$

we now have $u > dp^s$. Hence

$$m := \text{td}(p^s(d, d, d), (u, u, u)) = 3(u - dp^s) < 1.$$

Once again we get

$$\begin{aligned}
\delta^*(d, d, d) &= p^{-s}(1 - m) \\
&= p^{-s}(1 - 3u - 3dp^s) \\
&= p^n(1 - 3u) + 3d \\
&= -(3d - 1) + 3d \\
&= 1.
\end{aligned}$$

We conclude as above that A has the WLP. $\square$

As a corollary we obtain [7, Conjecture 3.9].

Corollary 2.7. *Let K be a field of characteristic 2. Then the Artinian complete intersection $A := K[X, Y, Z]/(X^d, Y^d, Z^d)$ has the weak Lefschetz property if and only if $d = \lfloor \frac{2^t+1}{3} \rfloor$ for some positive integer t .*

Proof. Let $n \in \mathbb{N}$ such that

$$\frac{3d}{2} > 2^n > \frac{3d}{4}$$

(note that there is only one such n since $\frac{3d}{4}$ is the half of $\frac{3d}{2}$). This n corresponds to $k = 0$ and is the exponent we have to consider by Theorem 2.6. So it follows from part (1) of Theorem 2.6 that the algebra A never enjoys the WLP for d even. So we may assume that d is odd. If

$$\frac{3d-1}{2} > 2^n > \frac{3d+1}{4}$$

holds then again A does not have the WLP. So A does have the WLP if either

$$\frac{3d-1}{2} \leq 2^n < \frac{3d}{2} \text{ or } \frac{3d}{2} < 2^{n+1} \leq \frac{3d+1}{2}$$

holds, i.e., if we have either $3d-1 = 2^{n+1}$ or $3d+1 = 2^{n+2}$. This gives the assertion of the corollary. $\square$

Remark 2.8. As remarked in [7] and indicated in our proof, Corollary 2.7 implies that the monomial complete intersection $K[X, Y, Z]/(X^d, Y^d, Z^d)$ does *not* have the WLP in characteristic 2 if d is even.

Theorem 2.6 implies in particular that for given d the weak Lefschetz property might only fail in characteristic $p \leq \frac{3}{2}d$. It is easy to generate the list of exceptional characteristics with the help of this numerical criterion.

Corollary 2.9. *Let d be odd and let p be a prime factor of d . Then the Artinian algebra $K[X, Y, Z]/(X^d, Y^d, Z^d)$ does not have the weak Lefschetz property in characteristic p .*

Proof. We write $d = p^n u$ with $u = 2k + 1$ odd, $n \geq 1$. Then

$$p^n = \frac{d}{2k+1} = \frac{3d}{6k+3}.$$

Since the numerator is larger than the denominator, this number is strictly between $\frac{3d+1}{6k+3+1}$ and $\frac{3d-1}{6k+3-1}$, so this fulfills the condition of Theorem 2.6(2). $\square$

Remark 2.8 and Corollary 2.9 imply that only for $d = 1$ the WLP holds in all characteristics. We will see in the examples below that for d even the weak Lefschetz property can hold in characteristics dividing d (but not in characteristic 2).

Example 2.10. We consider d even and determine the exceptional prime numbers (here we mean by *exceptional* that the Artinian complete intersection $A = K[X, Y, Z]/(X^d, Y^d, Z^d)$ does not enjoy the WLP in these characteristics).

$d = 2$. The only exceptional prime number is 2.

$d = 4$. The condition for $k = 0$ is $12/2 = 6 > p^n > 12/4 = 3$, hence the exceptional prime numbers are 2 and 5 (no larger k have to be considered).

$d = 6$. For $k = 0$ we get $9 > p^n > 4.5$, which yields the exceptional primes 2, 5, 7 (no larger k). The prime number 3 divides d , but the weak Lefschetz property does hold in characteristic 3.

$d = 8$. For $k = 0$ we get $12 > p^n > 6$, which yields the exceptional primes 2, 3, 7, 11 (no larger k).

$d = 10$. For $k = 0$ we get the exceptional primes 2, 3, 11, 13 (no larger k).

$d = 12$. For $k = 0$ we get the exceptional primes 2, 11, 13, 17 (no larger k).

$d = 14$. For $k = 0$ we get the condition $21 > p^n > 10.5$, which yields the exceptional primes 2, 11, 13, 17, 19. For $k = 1$ we get the condition $\frac{42}{8} > p^n > \frac{42}{10}$, which yields $p = 5$.

$d = 20$. For $k = 0$ we get the exceptional primes 2, 3, 5, 17, 19, 23, 29 and for $k = 1$ we also get 7. Note that 5 does divide d and the algebra does not have the weak Lefschetz property in characteristic 5.

Example 2.11. We consider d odd and determine the exceptional prime numbers.

$d = 1$. For $k = 0$ we get the condition $1 > p^n > 1$, which has no solution, hence $K[X, Y, Z]/(X, Y, Z) \cong K$ has the weak Lefschetz property in every characteristic, which is clear anyway.

$d = 3$. The condition for $k = 0$ is $8/2 = 4 > p^n > 10/4 = 2.5$, hence the only exceptional prime number is 3 (no larger k have to be considered).

$d = 5$. For $k = 0$ we get $7 > p^n > 4$, which yields the only exceptional prime 5 (no larger k). The prime number 7 fulfills $7 = \frac{14}{2} = \frac{3d-1}{2}$, which corresponds to the second case in the proof of Lemma 2.5. For $p = 7$ the Han number is $s = -1$, but the syzygy gap is 1 and not 3.

$d = 7$. For $k = 0$ we get $10 > p^n > 5.5$, which yields the exceptional primes 2, 3, 7 (no larger k).

$d = 9$. For $k = 0$ we get the exceptional primes 2, 3 and 11 (no larger k).

$d = 31$. For $k = 0$ we get the condition $46 > p^n > 23.5$, which yields the exceptional primes 2, 3, 5, 29, 31, 37, 41, 43. For $k = 1$ we get the condition $\frac{92}{8} > p^n > \frac{94}{10}$, which yields also $p = 11$.

REFERENCES

1. H. Brenner, *Looking out for stable syzygy bundles*, Adv. Math. **219** (2008), no. 2, 401–427.
2. H. Brenner and A. Kaid, *Syzygy bundles on $\mathbb{P}^2$ and the weak Lefschetz property*, Illinois J. Math. **51** (2007), no. 4, 1299–1308.
3. L. Ein, *Stable vector bundles on projective spaces in char p* , Math. Ann. **254** (1980), 53–72.
4. C. Han, *The Hilbert-Kunz function of a diagonal hypersurface*, PhD thesis, Brandeis University, 1991.
5. T. Harima, J. C. Migliore, U. Nagel, and J. Watanabe, *The weak and strong Lefschetz properties for Artinian k -algebras*, J. Algebra **262** (2003), 99–126.
6. D. Huybrechts and M. Lehn, *The Geometry of Moduli Spaces of Sheaves*, Viehweg, 1997.
7. J. Li and F. Zanello, *Monomial complete intersections, the weak Lefschetz property and plane partitions*, Preprint (2010), arXiv:1002.4400.
8. J. Migliore, R. M. Miró-Roig, and U. Nagel, *Monomial ideals, almost complete intersections and the weak Lefschetz property*, Preprint (2008), to appear in Trans. Amer. Math. Soc., arXiv:0811.1023.
9. P. Monsky, *Mason’s theorem and syzygy gaps*, J. Algebra **303** (2006), no. 1, 373–381.

UNIVERSITÄT OSNABRÜCK, FACHBEREICH 6: MATHEMATIK/INFORMATIK, ALBRECHTSTR. 28A, 49069 OSNABRÜCK, GERMANY

E-mail address: `hbrenner@uni-osnabrueck.de` and `akaid@uni-osnabrueck.de`